

# Gamma Knife Radiosurgery for AVM's: A Clinical Series of 199 Patients

Selcuk Peker

Koc University School of Medicine, Neurosurgery, Istanbul, Turkey

**Objective(s):** The aim of this study was to evaluate outcomes following gamma knife radiosurgery (GKRS) for arteriovenous malformations (AVMs).

**Methods:** The present series included 199 patients (3-74 years of age) who underwent GKRS for cerebral AVMs. In 43% of the patients the main symptom was hemorrhage. The median AVM volume was 4,9 cm<sup>3</sup>, and the median marginal dose was 22 Gy. All patients continued to have follow-up for more than 60 months with serial magnetic resonance images (MRIs)/angiograms.

**Results:** Follow-up radiological examinations confirmed complete obliteration in 140 patients (70%). Postradiosurgical hemorrhage was observed in 6 (3%) patients. Adverse radiation effects were seen in 3 patients (1.5%). The Spetzler-Martin grade, marginal dosis, volume and Pollock-Flickinger grade were found to be reliable predictors for total obliteration.

**Conclusion(s):** The results of the present study demonstrated a high total obliteration rate with low complications can be achievable in the GKRS treatment of AVM's.

Frontiers of Radiosurgery - October 24-25, 2019 - Milano, Italy